# The Diabetes Educator

# National Standards for Diabetes Self-Management Education and Support

Linda Haas, Melinda Maryniuk, Joni Beck, Carla E. Cox, Paulina Duker, Laura Edwards, Ed Fisher, Lenita Hanson, Daniel Kent, Leslie Kolb, Sue McLaughlin, Eric Orzeck, John D. Piette, Andrew S. Rhinehart, Russell Rothman, Sara Sklaroff, Donna Tomky and Gretchen Youssef The Diabetes Educator 2012 38: 619 DOI: 10.1177/0145721712455997

> The online version of this article can be found at: http://tde.sagepub.com/content/38/5/619

> > Published by: SAGE http://www.sagepublications.com

> > > On behalf of:



American Association of Diabetes Educators

Additional services and information for The Diabetes Educator can be found at:

Email Alerts: http://tde.sagepub.com/cgi/alerts

Subscriptions: http://tde.sagepub.com/subscriptions

Reprints: http://www.sagepub.com/journalsReprints.nav

Permissions: http://www.sagepub.com/journalsPermissions.nav

>> Version of Record - Sep 20, 2012

What is This?

# National Standards for **Diabetes Self-Management Education and Support**

By the most recent estimates, 18.8 million people in the United States have been diagnosed with diabetes, and an additional 7 million are believed to be living with undiagnosed diabetes. At the same time, 79 million people are estimated to have blood glucose levels in the prediabetes range. Thus, more than 100 million Americans are at risk of developing the devastating complications of diabetes.<sup>1</sup>

Diabetes self-management education (DSME) is a critical element of care for all people with diabetes and is necessary to prevent or delay the complications of diabetes.<sup>2-6</sup> Elements of DSME related to lifestyle change are also essential for people with prediabetes, as part of efforts to prevent the disease.<sup>7,8</sup> The National Standards for Diabetes Self-Management Education are designed to define quality DSME and support and to assist diabetes educators in providing evidence-based education and self-management support. The standards are applicable to educators in solo practice as well as those in large multicenter programs-and everyone in between. There are many good models for the provision of diabetes education and support. The standards do not endorse any one approach but rather seek to delineate the commonalities among effective and excellent self-management education strategies. These are the standards used in the field for recognition and accreditation. They also serve as a guide for nonaccredited and nonrecognized providers and programs.

Because of the dynamic nature of health care and diabetes-related research, the standards are reviewed and revised approximately every 5 years by key stakeholders and experts within the diabetes education community. In the fall of 2011, a task force was jointly convened by the Task Force Members:

Linda Haas, PhC, RN, CDE (Chair)
Melinda Maryniuk, MED, RD, CDE (Chair)
Joni Beck, PharmD, CDE, BC-ADM
Carla E. Cox, PhD, RD, CDE, CSSD
Paulina Duker, MPH, RN, BC-ADM, CDE
Laura Edwards, RN, MPA
Ed Fisher, PhD
enita Hanson, MD, CDE, FACE, FACP
Daniel Kent, PharmD, BS, CDE
eslie Kolb, RN, BSN, MBA
Sue McLaughlin, BS, RD, CDE, CPT
Eric Orzeck, MD, FACE, CDE
John D. Piette, PhD
Andrew S. Rhinehart, MD, FACP, CDE
Russell Rothman, MD, MPP
Sara Sklaroff
Donna Tomky, MSN, RN, C-NP, CDE, FAADE
Gretchen Youssef, MS, RD, CDE
/A Dugat Sound Haalth Care Sustem Hospital and Specialty Medicine, Seattle, Weshington, Joalin

re Syste alty Mec Diabetes Center, Boston, Massachusetts; University of Oklahoma Health Sciences Center, College of Medicine, Edmond, Oklahoma; Western Montana Clinic, Missoula, Montana; American Diabetes Association, Alexandria, Virginia; Center for Healthy North Carolina, Apex, North Carolina; Department of Health Behavior and Health Education, Gillings School of Global Public Health. University of North Carolina at Chapel Hill, Chapel Hill, North Carolina: Ultracare Endocrine and Diabetes Consultants, Venice, Florida: Group Health Central Specialty Clinic, Seattle, Washington: American Association of Diabetes Educators, Chicago, Illinois; On Site Health and Wellness, LLC, Omaha, Nebraska; Endocrinology Associates, Houston, Texas; Center for Global Health, VA Center for Clinical Management Research; University of Michigan Health System, Ann Arbor, Michigan; Johnston Memorial Diabetes Care Center, Abingdon, Virginia; Center for Health Services Research, Vanderbilt University Medical Center, Nashville, Tennessee; Writer and person with diabetes, Washington, DC; Department of Endocrinology and Diabetes, ABQ Health Partners, Albuquerque, New Mexico; MedStar Diabetes Institute, MedStar Health, Washington, DC

Correspondence to Leslie E. Kolb, RN, BSN, MBA, Director of Accreditation and Quality Initiatives, American Association of Diabetes Educators, 200 West Madison, Suite 800, Chicago, IL 60606 (lkolb@aadenet.org); and Paulina N. Duker, MPH, RN, BC-ADM, CDE, Vice President, Diabetes Education and Clinical Programs, American Diabetes Association, 1701 North Beauregard Street, Alexandria, VA 22311 (pduker@diabetes.org).

Acknowledgments: Linda Haas and Melinda Maryniuk were co-chairs of the Task Force. Leslie E. Kolb, RN, BSN, MBA, and Paulina N. Duker, MPH, RN, BC-ADM, worked on behalf of AADE and ADA, respectively, coordinating the effort on this document. Sara Sklaroff was the medical writer

DOI: 10.1177/0145721712455997

© 2012 The Author(s)

American Association of Diabetes Educators (AADE) and the American Diabetes Association. Members of the task force included experts from the areas of public health, underserved populations including rural primary care and other rural health services, individual practice, large urban specialty practice, and urban hospitals. They also included people with diabetes, diabetes researchers, certified diabetes educators, registered nurses, registered dietitians, physicians, pharmacists, and a psychologist. The task force was charged with reviewing the current National Standards for Diabetes Self-Management Education for their appropriateness, relevance, and scientific basis and updating them based on the available evidence and expert consensus.

The task force made the decision to change the name of the standards from the National Standards for Diabetes Self-Management Education to the National Standards for Diabetes Self-Management Education and Support. This name change is intended to codify the significance of ongoing support for people with diabetes, particularly to encourage behavior change and the maintenance of healthy diabetes-related behaviors and to address psychosocial concerns. Given that self-management does not stop when a patient leaves the educator's office, selfmanagement support must be an ongoing process.

Although the term *diabetes* is used predominantly, the standards should be understood to apply to the education and support of people with prediabetes. Currently, there are significant barriers to the provision of education and support to those with prediabetes. And yet, the strategies for supporting successful behavior change and the healthy behaviors recommended for people with prediabetes are largely identical to those for people with diabetes. As barriers to care are overcome, providers of DSME and diabetes self-management support (DSMS), given their training and experience, are particularly well equipped to assist people with prediabetes in developing and maintaining behaviors that can prevent or delay the onset of diabetes.

Many people with diabetes have or are at risk for developing comorbidities, including heart disease, lipid abnormalities, nerve damage, hypertension, and depression. In addition, the diagnosis, progression, and daily work of managing the disease can take a major emotional toll on people with diabetes that makes self-care even more difficult.<sup>9</sup> The standards encourage providers of DSME and DSMS to address the entire panorama of each participant's clinical profile. Regular communication among the members of participants' health care teams is essential to ensure high-quality, effective education and support for people with diabetes and prediabetes.

In the course of its work on the standards, the task force identified areas in which there is currently an insufficient amount of research. In particular, there are 3 areas for which the task force recommends additional research:

- 1. What is the influence of organizational structure on the effectiveness of the provision of DSME?
- 2. What is the impact of using a structured curriculum in DSME?
- 3. What training should be required for those community, lay, or peer workers without training in health or diabetes who are to participate in the provision of DSME and provide DSMS?

Finally, the standards emphasize that the person with diabetes is at the center of the entire diabetes education and support process. It is people with diabetes who do the hard work of managing their condition, day in and day out. The educator's role, first and foremost, is to make that work easier.<sup>10</sup>

# Definitions

Diabetes self-management education: the ongoing process of facilitating the knowledge, skill, and ability necessary for prediabetes and diabetes self-care. This process incorporates the needs, goals, and life experiences of the person with diabetes or prediabetes and is guided by evidence-based standards. The overall objectives of DSME are to support informed decision making, self-care behaviors, problem solving, and active collaboration with the health care team and to improve clinical outcomes, health status, and quality of life.

*Diabetes self-management support*: activities that assist the person with prediabetes or diabetes in implementing and sustaining the behaviors needed to manage his or her condition on an ongoing basis beyond or outside of formal self-management training. The type of support provided can be behavioral, educational, psychosocial, or clinical.<sup>11-15</sup>

# Standard 1

#### **Internal Structure**

The provider(s) of DSME will document an organizational structure, mission statement, and goals. For those providers working within a larger organization, that organization will recognize and support quality DSME as an integral component of diabetes care.

Documentation of an organizational structure, mission statement, and goals can lead to efficient and effective provision of DSME and DSMS. In the business literature, case studies and case report investigations of successful management strategies emphasize the importance of clear goals and objectives, defined relationships and roles, and managerial support. Business and health policy experts and organizations emphasize written commitments, policies, support, and the importance of outcomes reporting to maintain ongoing support or commitment.<sup>16,17</sup>

Documentation of an organizational structure that delineates channels of communication and represents institutional commitment to the educational entity is critical for success. According to the Joint Commission on Accreditation of Healthcare Organizations, this type of documentation is equally important for small and large health care organizations.<sup>18</sup> Health care and business experts overwhelmingly agree that documentation of the process of providing services is a critical factor in clear communication and provides a solid basis from which to deliver quality diabetes education. In 2010, the joint commission published the Disease-Specific Care Certification Manual, which outlines standards and performance measurements for chronic care programs and disease management services, including "supporting self-management."18

# Standard 2

# **External Input**

The provider(s) of DSME will seek ongoing input from external stakeholders and experts to promote program quality.

For individual and group providers of DSME and DSMS, external input is vital to maintain an up-to-date, effective program. Broad participation of community stakeholders, including people with diabetes, health professionals, and community interest groups, will increase the program's knowledge of the local population and allow the provider to better serve the community. Often, but not always, this external input is best achieved by the establishment of a formal advisory board. The DSME and DSMS provider(s) must have a documented plan for seeking outside input and acting on it. The goal of external input and discussion in the program planning process is to foster ideas that will enhance the quality of the DSME and/or DSMS being provided while building bridges to key stakeholders.<sup>19</sup> The result is effective, dynamic DSME that is patient centered, more responsive to consumer-identified needs and the needs of the community, more culturally relevant, and more appealing to consumers.<sup>17,19,20</sup>

# Standard 3

# Access

The provider(s) of DSME will determine whom to serve, how best to deliver diabetes education to that population, and what resources can provide ongoing support for that population.

Currently, the majority of people with diabetes and prediabetes do not receive any structured diabetes education.<sup>19,20</sup> While there are many barriers to DSME, one crucial issue is access.<sup>21</sup> Providers of DSME can help address this issue by doing the following.

\* Clarifying the specific population to be served.

Understanding the community, service area, or regional demographics is crucial to ensuring that as many people as possible are being reached, including those who do not frequently attend clinical appointments.<sup>9,17,22-24</sup>

\* Determining that population's self-management education and support needs.

Different individuals, their families, and communities need different types of education and support.<sup>25</sup> The provider of DSME needs to work to ensure that the necessary education alternatives are available.<sup>25-27</sup> This means understanding the population's demographic characteristics, such as ethnic/cultural background, gender, and age, as well as their levels of formal education, literacy, and numeracy.<sup>28-31</sup> It may also entail identifying resources outside the provider's practice that can assist in the ongoing support of the participant.

\* Identifying access issues and working to overcome them.

It is essential to determine factors that prevent people with diabetes from receiving self-management education and support. The assessment process includes the identification of these barriers to access.<sup>32-34</sup> These barriers may

include the socioeconomic or cultural factors mentioned above, as well as, for example, health insurance shortfalls and the lack of encouragement from other health providers to encourage their patients to pursue diabetes education.<sup>35,36</sup>

# **Standard 4**

## **Program Coordination**

A coordinator will be designated to oversee the DSME program. The coordinator will have oversight responsibility for the planning, implementation, and evaluation of education services.

Coordination is essential to ensure that quality DSME and support are delivered through an organized, systematic process.<sup>37,38</sup> As the field of DSME continues to evolve, the coordinator plays a pivotal role in ensuring accountability and continuity in the education program.<sup>39-41</sup> The coordinator's role may be viewed as that of coordinating the program (or education process) and/or supporting the coordination of the many aspects of selfmanagement in the continuum of diabetes and related conditions when feasible.<sup>42-49</sup> This oversight includes designing an education program or service that helps the participant access needed resources and assists him or her in navigating the health care system.<sup>37,50-55</sup>

The individual serving as the coordinator will have knowledge of the lifelong process of managing a chronic disease and facilitating behavior change, in addition to experience with program and/or clinical management.<sup>56-59</sup> In some cases, particularly solo or other small practices, the coordinator may also provide DSME and/or DSMS.

# Standard 5

# Instructional Staff

One or more instructors will provide DSME and, when applicable, DSMS. At least one of the instructors responsible for designing and planning DSME and DSMS will be an RN, RD, or pharmacist with training and experience pertinent to DSME, or another professional with certification in diabetes care and education, such as a CDE or BC-ADM. Other health workers can contribute to DSME and provide DSMS with appropriate training in diabetes and with supervision and support.

Historically, nurses and dietitians were the main providers of diabetes education.<sup>3,4,60-64</sup> In recent years, the role of the diabetes educator has expanded to other disciplines, particularly pharmacists.<sup>65-67</sup> Reviews comparing the effectiveness of different disciplines for education have not identified clear differences in the quality of services delivered by different professionals.<sup>3-5</sup> However, the literature favors the registered nurse, registered dietitian, and pharmacist serving both as the key primary instructors for diabetes education and as members of the multidisciplinary team responsible for designing the curriculum and assisting in the delivery of DSME.<sup>1-7</sup> Expert consensus supports the need for specialized diabetes and educational training beyond academic preparation for the primary instructors on the diabetes team.<sup>68-71</sup> Professionals serving as instructors must document appropriate continuing education or comparable activities to ensure their continuing competence to serve in their instructional, training, and oversight roles.<sup>72</sup>

Reflecting the evolving health care environment, a number of studies have endorsed a multidisciplinary team approach to diabetes care, education, and support. The disciplines that may be involved include, but are not limited to, physicians, psychologists and other mental health specialists, physical activity specialists (including physical therapists, occupational therapists, and exercise physiologists), optometrists, and podiatrists.73-75 More recently, health educators (e.g. certified health education specialists and certified medical assistants), case managers, lay health and community workers,<sup>76-83</sup> and peer counselors or educators<sup>84,85</sup> have been shown to contribute effectively as part of the DSME team and in providing DSMS. While DSME and DSMS are often provided within the framework of a collaborative and integrated team approach, it is crucial that the individual with diabetes be viewed as central to the team and that he or she take an active role.

Certification as a diabetes educator by the National Certification Board for Diabetes Educators is one way that a health professional can demonstrate mastery of a specific body of knowledge, and this certification has become an accepted credential in the diabetes community.<sup>86</sup> An additional credential that indicates specialized training beyond basic preparation is board certification in advanced diabetes management offered by the AADE, which is available for nurses, dietitians, pharmacists, physicians, and physician assistants.<sup>73,75,87</sup>

Individuals who serve as lay health and community workers and peer counselors or educators may contribute to the provision of DSME instruction and provide DSMS if they have received training in diabetes management,

Volume 38, Number 5, September/October 2012

the teaching of self-management skills, group facilitation, and emotional support. For these individuals, a system must be in place that ensures supervision of the services they provide by a diabetes educator or other health care professional and professional backup to address clinical problems or questions beyond their training.<sup>88-90</sup>

For services outside the expertise of any provider of DSME and DSMS, a mechanism must be in place to ensure that the individual with diabetes is connected with appropriately trained and credentialed providers.

# **Standard 6**

# Curriculum

A written curriculum reflecting current evidence and practice guidelines, with criteria for evaluating outcomes, will serve as the framework for the provision of DSME. The needs of the individual participant will determine which parts of the curriculum will be provided to that individual.

People with prediabetes and diabetes and their families and caregivers have much to learn to become effective self-managers of their condition. DSME can provide this education via an up-to-date, evidence-based, and flexible curriculum.<sup>8,91</sup>

The curriculum is a coordinated set of courses and educational experiences. It also specifies learning outcomes and effective teaching strategies.<sup>92,93</sup> The curriculum must be dynamic and reflect current evidence and practice guidelines.<sup>93-97</sup> Recent education research endorses the inclusion of practical, problem-solving approaches, collaborative care, psychosocial issues, behavior change, and strategies to sustain self-management efforts.<sup>12,13,19,73,86,98-101</sup>

The following core topics are commonly part of the curriculum taught in comprehensive programs that have demonstrated successful outcomes.<sup>2,3,5,91,102-104</sup>

- Describing the diabetes disease process and treatment options
- Incorporating nutritional management into lifestyle
- Incorporating physical activity into lifestyle
- Using medication safely and for maximum therapeutic effectiveness
- Monitoring blood glucose and other parameters and interpreting and using the results for self-management decision making
- Preventing, detecting, and treating acute complications

- Preventing, detecting, and treating chronic complications
- Developing personal strategies to address psychosocial issues and concerns
- Developing personal strategies to promote health and behavior change

While the content areas listed above provide a solid outline for a diabetes education and support curriculum, it is crucial that the content be tailored to match each individual's needs and be adapted as necessary for age, type of diabetes (including prediabetes and diabetes in pregnancy), cultural factors, health literacy and numeracy, and comorbidities.<sup>14,105-108</sup> The content areas will be able to be adapted for all practice settings.

Approaches to education that are interactive and patient centered have been shown to be effective.<sup>12,13,109-112</sup> Also crucial is the development of action-oriented behavioral goals and objectives.<sup>12-14,113</sup> Creative, patient-centered, experience-based delivery methods—beyond the mere acquisition of knowledge—are effective for supporting informed decision making and meaningful behavior change and addressing psychosocial concerns.<sup>114,115</sup>

# Standard 7

# Individualization

The diabetes self-management, education, and support needs of each participant will be assessed by one or more instructors. The participant and instructor(s) will then together develop an individualized education and support plan focused on behavior change.

Research has demonstrated the importance of individualizing diabetes education to each participant's needs.<sup>116</sup> The assessment process is used to identify what those needs are and to facilitate the selection of appropriate educational and behavioral interventions and self-management support strategies, guided by evidence.<sup>2,63,116-118</sup> The assessment must garner information about the individual's medical history, age, cultural influences, health beliefs and attitudes, diabetes knowledge, diabetes self-management skills and behaviors, emotional response to diabetes, readiness to learn, literacy level (including health literacy and numeracy), physical limitations, family support, and financial status.<sup>11,106,108,117,119-128</sup>

The education and support plan that the participant and instructors develop will be rooted in evidence-based approaches to effective health communication and

education while taking into consideration participant barriers, abilities, and expectations. The instructor will employ clear health communication principles, avoiding jargon, making information culturally relevant, using language- and literacy-appropriate education materials, and using interpreter services when indicated.<sup>107,129,130,131</sup> Evidence-based communication strategies are also effective, such as collaborative goal setting, motivational interviewing, cognitive behavior change strategies, problem solving, self-efficacy enhancement, and relapse prevention strategies.<sup>101,132-134</sup> Periodic reassessment can determine whether there is need for additional or different interventions and future reassessment.6,71,134-137 A variety of assessment modalities, including telephone follow-up and other information technologies (eg, Internet, text messaging, automated phone calls), may augment face-to-face assessments.71,87,138-141

The assessment and education plan, intervention, and outcomes will be documented in the education/health record. Documentation of participant encounters will guide the education process, provide evidence of communication among instructional staff and other members of the participant's health care team, prevent duplication of services, and demonstrate adherence to guidelines.<sup>117,135,142,143</sup> Providing information to other members of the participant's health care team through documentation of educational objectives and personal behavioral goals increases the likelihood that all members will work in collaboration.<sup>86,143</sup> Evidence suggests that the development of standardized procedures for documentation, training health professionals to document appropriately, and the use of structured standardized forms based on current practice guidelines can improve documentation and may ultimately improve quality of care.135,143-145

# Standard 8

# **Ongoing Support**

The participant and instructor(s) will together develop a personalized follow-up plan for ongoing selfmanagement support. The participant's outcomes and goals and the plan for ongoing self-management support will be communicated to other members of the healthcare team.

While DSME is necessary and effective, it does not in itself guarantee a lifetime of effective diabetes self-care.<sup>113</sup> Initial improvements in participants' metabolic

and other outcomes have been found to diminish after approximately 6 months.<sup>3</sup> To sustain the level of selfmanagement needed to effectively manage prediabetes and diabetes over the long term, most participants need ongoing DSMS.<sup>15</sup>

The type of support provided can be behavioral, educational, psychosocial, or clinical.<sup>11-14</sup> A variety of strategies are available for providing DSMS both within and outside the DSME organization. Some patients benefit from working with a nurse case manager.<sup>6,86,146</sup> Case management for DSMS can include reminders about needed follow-up care and tests, medication management, education, behavioral goal setting, psychosocial support, and connection to community resources.

The effectiveness of providing DSMS through disease management programs, trained peers and community health workers, community-based programs, information technology, ongoing education, support groups, and medical nutrition therapy has also been established.<sup>7-11,88-90,86,142</sup>

While the primary responsibility for diabetes education belongs to the providers of DSME, participants benefit by receiving reinforcement of content and behavioral goals from their entire health care team.<sup>135</sup> Additionally, many patients receive DSMS through their primary care provider. Thus, communication among the team regarding the patient's educational outcomes, goals, and DSMS plan is essential to ensure that people with diabetes receive support that meets their needs and is reinforced and consistent among the health care team members.

Because self-management takes place in participants' daily lives and not in clinical or educational settings, patients will be assisted to formulate a plan to find community-based resources that may support their ongoing diabetes self-management. Ideally, DSME and DSMS providers will work with participants to identify such services and, when possible, track those that have been effective with patients, while communicating with providers of community-based resources to better integrate them into patients' overall care and ongoing support.

# Standard 9

## **Patient Progress**

The provider(s) of DSME and DSMS will monitor whether participants are achieving their personal diabetes selfmanagement goals and other outcome(s) as a way to evaluate the effectiveness of the educational intervention(s), using appropriate measurement techniques.

Volume 38, Number 5, September/October 2012

Effective diabetes self-management can be a significant contributor to long-term, positive health outcomes. The provider(s) of DSME and DSMS will assess each participant's personal self-management goals and his or her progress toward those goals.<sup>147,148</sup>

The AADE Outcome Standards for Diabetes Education specify behavior change as the key outcome and provide a useful framework for assessment and documentation. The AADE7<sup>TM</sup> lists 7 essential factors: healthy eating, physical activity, taking medications, monitoring, diabetes self-care related problem solving, reducing risks of acute and chronic complications, and psychosocial aspects of living with diabetes.<sup>93,149,150</sup> Differences in behaviors, health beliefs, and culture as well as their emotional response to diabetes can have a significant impact on how participants understand their illness and engage in self-management. DSME providers who account for these differences when collaborating with participants on the design of personalized DSME or DSMS programs can improve participant outcomes.<sup>151,152</sup>

Assessments of participant outcomes must occur at appropriate intervals. The interval depends on the nature of the outcome itself and the time frame specified based on the participant's personal goals. For some areas, the indicators, measures, and time frames will be based on guidelines from professional organizations or government agencies.

# Standard 10

# **Quality Improvement**

The provider(s) of DSME will measure the effectiveness of the education and support and look for ways to improve any identified gaps in services or service quality, using a systematic review of process and outcome data.

Diabetes education must be responsive to advances in knowledge, treatment strategies, education strategies, and psychosocial interventions, as well as consumer trends and the changing health care environment. By measuring and monitoring both process and outcome data on an ongoing basis, providers of DSME can identify areas of improvement and make adjustments in participant engagement strategies and program offerings accordingly.

The Institute for Healthcare Improvement suggests 3 fundamental questions that should be answered by an improvement process:<sup>153</sup>

- \* What are we trying to accomplish?
- \* How will we know a change is an improvement?
- \* What changes can we make that will result in an improvement?

Once areas for improvement are identified, the DSME provider must designate timelines and important milestones, including data collection, analysis, and presentation of results.<sup>154</sup> Measuring processes and outcomes helps to ensure that change is successful without causing additional problems in the system. Outcome measures indicate the result of a process (ie, whether changes are actually leading to improvement), while process measures provide information about what caused those results.<sup>154</sup> Process measures are often targeted to those processes that typically affect the most important outcomes.

# References

- 1. Centers for Disease Control and Prevention. *National Diabetes Fact Sheet: National Estimates and General Information on Diabetes and Prediabetes in the United States, 2011.* Atlanta, GA: Centers for Disease Control and Prevention; 2011.
- 2. Brown SA. Interventions to promote diabetes self-management: state of the science. *Diabetes Educ.* 1999;25(6)(suppl):52-61.
- 3. Norris SL, Lau J, Smith SJ, et al. Self-management education for adults with type 2 diabetes: a meta-analysis of the effect on glycemic control. *Diabetes Care*. 2002;25(7):1159-1171.
- 4. Gary TL, Genkinger JM, Guallar E, et al. Meta-analysis of randomized educational and behavioral interventions in type 2 diabetes. *Diabetes Educ.* 2003;29(3):488-501.
- Deakin T, McShane CE, Cade JE, et al. Group based training for self-management strategies in people with type 2 diabetes mellitus. *Cochrane Database Syst Rev.* 2005;2:CD003417.
- Renders CM, Valk GD, Griffin SJ, et al. Interventions to improve the management of diabetes in primary care, outpatient, and community settings: a systematic review. *Diabetes Care*. 2001;24(10):1821-1833.
- 7. Ratner RE. An update on the Diabetes Prevention Program. *Endocr Pract*. 2006;12(suppl 1):20-24.
- 8. Diabetes Prevention Program Research Group. The Diabetes Prevention Program (DPP): description of lifestyle intervention. *Diabetes Care*. 2002;25(12):2165-2171.
- Peyrot M, Rubin RR, Funnell MM, et al. Access to diabetes selfmanagement education: results of national surveys of patients, educators, and physicians. *Diabetes Educ.* 2009;35(2):246-248, 252-246, 258-263.
- Inzucchi SE, Bergenstal RM, Buse JB, et al. Management of hyperglycemia in type 2 diabetes: a patient-centered approach. Position statement of the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetes Care*. 2012;35(6):1364-1379.
- 11. Anderson RM, Funnell MM, Nwankwo R, et al. Evaluating a problem-based empowerment program for African Americans with diabetes: results of a randomized controlled trial. *Ethn Dis.* 2005;15(4):671-678.

- Tang TS, Gillard ML, Funnell MM, et al. Developing a new generation of ongoing: diabetes self-management support interventions. A preliminary report. *Diabetes Educ.* 2005;31(1):91-97.
- Funnell MM, Nwankwo R, Gillard ML, et al. Implementing an empowerment-based diabetes self-management education program. *Diabetes Educ*. 2005;31(1):53, 55-56, 61.
- Glazier RH, Bajcar J, Kennie NR, et al. A systematic review of interventions to improve diabetes care in socially disadvantaged populations. *Diabetes Care*. 2006;29(7):1675-1688.
- Fjeldsoe BS, Marshall AL, Miller YD. Behavior change interventions delivered by mobile telephone short-message service. *Am J Prev Med.* 2009;36(2):165-173.
- Armstrong G, Headrick L, Madigosky W, et al. Designing education to improve care. *Jt Comm J Qual Patient Saf.* 2012;38(1):5-14.
- Martin AL. Changes and consistencies in diabetes education over 5 years: results of the 2010 National Diabetes Education Practice Survey. *Diabetes Educ.* 2012;38(1):35-46.
- Joint Commission on Accreditation of Healthcare Organizations. Disease-Specific Care Certification Manual. Oakbrook Terrace, IL: Joint Commission on Accreditation of Healthcare Organizations; 2010.
- Siminerio LM, Piatt GA, Emerson S, et al. Deploying the chronic care model to implement and sustain diabetes self-management training programs. *Diabetes Educ*. 2006;32(2):253-260.
- Siminerio LM, Piatt G, Zgibor JC. Implementing the chronic care model for improvements in diabetes care and education in a rural primary care practice. *Diabetes Educ.* 2005;31(2):225-234.
- Boren SA, Fitzner KA, Panhalkar PS, et al. Costs and benefits associated with diabetes education: a review of the literature. *Diabetes Educ.* 2009;35(1):72-96.
- McWilliams JM, Meara E, Zaslavsky AM, et al. Health of previously uninsured adults after acquiring Medicare coverage. *JAMA*. 2007;298(24):2886-2894.
- Bell RA, Mayer-Davis EJ, Beyer JW, et al. Diabetes in non-Hispanic white youth: prevalence, incidence, and clinical characteristics. The SEARCH for Diabetes in Youth Study. *Diabetes Care*. 2009;32(suppl 2):S102-S111.
- Glasgow RE. Interactive media for diabetes self-management: issues in maximizing public health impact. *Med Decis Making*. 2010;30(6):745-758.
- Lorig K, Ritter PL, Villa FJ, et al. Community-based peer-led diabetes self-management: a randomized trial. *Diabetes Educ*. 2009;35(4):641-651.
- Duke SA, Colagiuri S, Colagiuri R. Individual patient education for people with type 2 diabetes mellitus. *Cochrane Database Syst Rev.* 2009;1:CD005268.
- Siminerio LM, Drab SR, Gabbay RA, et al. Diabetes educators: implementing the chronic care model. *Diabetes Educ*. 2008;34(3):451-456.
- Rosal MC, Ockene IS, Restrepo A, et al. Randomized trial of a literacy-sensitive, culturally tailored diabetes self-management intervention for low-income latinos: Latinos en control. *Diabetes Care*. 2011;34(4):838-844.
- Mayer-Davis EJ, Beyer J, Bell RA, et al. Diabetes in African American youth: prevalence, incidence, and clinical characteristics. The SEARCH for Diabetes in Youth Study. *Diabetes Care*. 2009;32(suppl 2):S112-S122.

- 30. Liu LL, Yi JP, Beyer J, et al. Type 1 and type 2 diabetes in Asian and Pacific Islander US youth: the SEARCH for Diabetes in Youth Study. *Diabetes Care*. 2009;32(suppl 2):S133-S140.
- Hill-Briggs F, Batts-Turner M, Gary TL, et al. Training community health workers as diabetes educators for urban African Americans: value added using participatory methods. *Prog Community Health Partnersh*. 2007;1(2):185-194.
- Unutzer J, Schoenbaum M, Katon WJ, et al. Healthcare costs associated with depression in medically ill fee-for-service Medicare participants. *J Am Geriatr Soc.* 2009;57(3):506-510.
- 33. Walker EA, Shmukler C, Ullman R, et al. Results of a successful telephonic intervention to improve diabetes control in urban adults: a randomized trial. *Diabetes Care*. 2011;34(1):2-7.
- Wubben DP, Vivian EM. Effects of pharmacist outpatient interventions on adults with diabetes mellitus: a systematic review. *Pharmacotherapy*. 2008;28(4):421-436.
- 35. Remler DK, Teresi JA, Weinstock RS, et al. Health care utilization and self-care behaviors of Medicare beneficiaries with diabetes: comparison of national and ethnically diverse underserved populations. *Popul Health Manag.* 2011;14(1):11-20.
- Peikes D, Chen A, Schore J, et al. Effects of care coordination on hospitalization, quality of care, and health care expenditures among Medicare beneficiaries: 15 randomized trials. *JAMA*. 2009;301(6):603-618.
- 37. Rothman RL, Malone R, Bryant B, et al. A randomized trial of a primary care-based disease management program to improve cardiovascular risk factors and glycated hemoglobin levels in patients with diabetes. *Am J Med.* 2005;118(3):276-284.
- Holmes-Walker DJ, Llewellyn AC, Farrell K. A transition care programme which improves diabetes control and reduces hospital admission rates in young adults with type 1 diabetes aged 15-25 years. *Diabet Med.* 2007;24(7):764-769.
- Glasgow RE, Nelson CC, Strycker LA, et al. Using RE-AIM metrics to evaluate diabetes self-management support interventions. *Am J Prev Med.* 2006;30(1):67-73.
- 40. Baker LC, Johnson SJ, Macaulay D, et al. Integrated telehealth and care management program for Medicare beneficiaries with chronic disease linked to savings. *Health Aff (Millwood)*. 2011;30(9):1689-1697.
- 41. Piatt GA, Anderson RM, Brooks MM, et al. 3-year follow-up of clinical and behavioral improvements following a multifaceted diabetes care intervention: results of a randomized controlled trial. *Diabetes Educ*. 2010;36(2):301-309.
- 42. Kerr EA, Heisler M, Krein SL, et al. Beyond comorbidity counts: how do comorbidity type and severity influence diabetes patients' treatment priorities and self-management? J Gen Intern Med. 2007;22(12):1635-1640.
- Bowen ME, Rothman RL. Multidisciplinary management of type 2 diabetes in children and adolescents. *J Multidiscip Healthc*. 2010;3:113-124.
- 44. Dejesus RS, Vickers KS, Stroebel RJ, et al. Primary care patient and provider preferences for diabetes care managers. *Patient Prefer Adherence*. 2010;4:181-186.
- 45. Stuckey HL, Dellasega C, Graber NJ, et al. Diabetes nurse case management and motivational interviewing for change (DYNAMIC): study design and baseline characteristics in the chronic care model for type 2 diabetes. *Contemp Clin Trials*. 2009;30(4):366-374.

Volume 38, Number 5, September/October 2012

- Heuer LJ, Hess C, Batson A. Cluster clinics for migrant Hispanic farmworkers with diabetes: perceptions, successes, and challenges. *Rural Remote Health*. 2006;6(1):469.
- 47. Cebul RD, Love TE, Jain AK, et al. Electronic health records and quality of diabetes care. *N Engl J Med.* 2011;365(9):825-833.
- 48. Rosal MC, White MJ, Borg A, et al. Translational research at community health centers: challenges and successes in recruiting and retaining low-income Latino patients with type 2 diabetes into a randomized clinical trial. *Diabetes Educ*. 2010;36(5):733-749.
- Austin SA, Claiborne N. Faith wellness collaboration: a community-based approach to address type II diabetes disparities in an African-American community. *Soc Work Health Care*. 2011;50(5):360-375.
- Parekh AK, Goodman RA, Gordon C, et al. Managing multiple chronic conditions: a strategic framework for improving health outcomes and quality of life. *Public Health Rep.* 2011;126(4):460-471.
- Rothman RL, So SA, Shin J, et al. Labor characteristics and program costs of a successful diabetes disease management program. *Am J Manag Care*. 2006;12(5):277-283.
- 52. May CR, Finch TL, Cornford J, et al. Integrating telecare for chronic disease management in the community: what needs to be done? *BMC Health Serv Res.* 2011;11:131.
- 53. Williams AS. Making diabetes education accessible for people with visual impairment. *Diabetes Educ.* 2009;35(4):612-621.
- Reichard A, Stolzle H. Diabetes among adults with cognitive limitations compared to individuals with no cognitive disabilities. *Intellect Dev Disabil*. 2011;49(3):141-154.
- Gimpel N, Marcee A, Kennedy K, et al. Patient perceptions of a community-based care coordination system. *Health Promot Pract*. 2010;11(2):173-181.
- 56. Welch G, Allen NA, Zagarins SE, et al. Comprehensive diabetes management program for poorly controlled Hispanic type 2 patients at a community health center. *Diabetes Educ*. 2011;37(5):680-688.
- Peterson KA, Radosevich DM, O'Connor PJ, et al. Improving diabetes care in practice: findings from the TRANSLATE trial. *Diabetes Care*. 2008;31(12):2238-2243.
- Bojadzievski T, Gabbay RA. Patient-centered medical home and diabetes. *Diabetes Care*. 2011;34(4):1047-1053.
- Wagner EH. The role of patient care teams in chronic disease management. *BMJ*. 2000;320(7234):569-572.
- 60. Koproski J, Pretto Z, Poretsky L. Effects of an intervention by a diabetes team in hospitalized patients with diabetes. *Diabetes Care*. 1997;20(10):1553-1555.
- Weinberger M, Kirkman MS, Samsa GP, et al. A nurse-coordinated intervention for primary care patients with non-insulin-dependent diabetes mellitus: impact on glycemic control and health-related quality of life. J Gen Intern Med. 1995;10(2):59-66.
- Spellbring AM. Nursing's role in health promotion. An overview. Nurs Clin North Am. 1991;26(4):805-814.
- 63. Glasgow RE, Toobert DJ, Hampson SE, et al. Improving self-care among older patients with type II diabetes: the "Sixty Something ..." Study. *Patient Educ Couns.* 1992;19(1):61-74.
- Delahanty L, Simkins SW, Camelon K. Expanded role of the dietitian in the Diabetes Control and Complications Trial: implications for clinical practice. The DCCT Research Group. J Am Diet Assoc. 1993;93(7):758-764, 767.
- 65. Cranor CW, Bunting BA, Christensen DB. The Asheville Project: long-term clinical and economic outcomes of a community

pharmacy diabetes care program. J Am Pharm Assoc (Wash). 2003;43(2):173-184.

- Garrett DG, Bluml BM. Patient self-management program for diabetes: first-year clinical, humanistic, and economic outcomes. *J Am Pharm Assoc (2003)*. 2005;45(2):130-137.
- Shane-McWhorter L, Fermo JD, Bultemeier NC, et al. National survey of pharmacist certified diabetes educators. *Pharmacotherapy*. 2002;22(12):1579-1593.
- Anderson RM, Donnelly MB, Dedrick RF, et al. The attitudes of nurses, dietitians, and physicians toward diabetes. *Diabetes Educ*. 1991;17(4):261-268.
- 69. Lorenz RA, Bubb J, Davis D, et al. Changing behavior: practical lessons from the diabetes control and complications trial. *Diabetes Care*. 1996;19(6):648-652.
- Ockene JK, Ockene IS, Quirk ME, et al. Physician training for patient-centered nutrition counseling in a lipid intervention trial. *Prev Med.* 1995;24(6):563-570.
- Leggett-Frazier N, Swanson MS, Vincent PA, et al. Telephone communications between diabetes clients and nurse educators. *Diabetes Educ.* 1997;23(3):287-293.
- 72. Baksi AK, Al-Mrayat M, Hogan D, et al. Peer advisers compared with specialist health professionals in delivering a training programme on self-management to people with diabetes: a randomized controlled trial. *Diabet Med.* 2008;25(9):1076-1082.
- Piatt GA, Orchard TJ, Emerson S, et al. Translating the chronic care model into the community: results from a randomized controlled trial of a multifaceted diabetes care intervention. *Diabetes Care*. 2006;29(4):811-817.
- Campbell EM, Redman S, Moffitt PS, et al. The relative effectiveness of educational and behavioral instruction programs for patients with NIDDM: a randomized trial. *Diabetes Educ*. 1996;22(4):379-386.
- 75. Emerson S. Implementing diabetes self-management education in primary care. *Diabetes Spectrum*. 2006;19(2):79-83.
- 76. Satterfield D, Burd C, Valdez L, et al. The "in-between people": participation of community health representatives and lay health workers in diabetes prevention and care in American Indian and Alaska Native communities. *Health Promot Pract.* 2002;3:66-175.
- 77. American Association of Diabetes Educators. Community Health Workers position statement. http://www.diabeteseducator.org/ ProfessionalResources/position/position\_statements.html. Accessed June 26, 2012.
- American Public Health Association. Support for community health workers to increase health access and to reduce health inequities. http://www.apha.org/advocacy/policy/policysearch/ default.htm?id=1393. Accessed June 26, 2012.
- Norris SL, Chowdhury FM, Van Le K, et al. Effectiveness of community health workers in the care of persons with diabetes. *Diabet Med.* 2006;23(5):544-556.
- Lewin SA, Dick J, Pond P, et al. Lay health workers in primary and community health care. *Cochrane Database Syst Rev.* 2005(1):CD004015.
- Lorig KR, Ritter P, Stewart AL, et al. Chronic disease self-management program: 2-year health status and health care utilization outcomes. *Med Care*. 2001;39(11):1217-1223.
- 82. Ruggiero L, Moadsiri A, Butler P, et al. Supporting diabetes selfcare in underserved populations: a randomized pilot study using medical assistant coaches. *Diabetes Educ*. 2010;36(1):127-131.

- 83. Spencer MS, Rosland AM, Kieffer EC, et al. Effectiveness of a community health worker intervention among African American and Latino adults with type 2 diabetes: a randomized controlled trial. Am J Public Health. 2011;101(12):2253-2260.
- Heisler M. Building Peer Support Programs to Manage Chronic Disease: Seven Models for Success. Oakland, CA: California Health Care Foundation; 2006.
- Long JA, Jahnle EC, Richardson DM, et al. Peer mentoring and financial incentives to improve glucose control in African American veterans: a randomized trial. *Ann Intern Med.* 2012;156(6):416-424.
- American Association of Diabetes Educators. The scope of practice, standards of practice, and standards of professional performance for diabetes educators. http://www.diabeteseducator.org/ DiabetesEducation/position/Scope\_x\_Standards.html. Accessed June 26, 2012.
- Valentine V, Kulkarni K, Hinnen D. Evolving roles: from diabetes educators to advanced diabetes managers. *Diabetes Educ*. 2003;29(4):598-602, 604, 606.
- American Association of Diabetes Educators. AADE guidelines for the practice of diabetes self-management education and training (DSME/T). *Diabetes Educ.* 2009;35(3):85S-107S.
- American Association of Diabetes Educators. Competencies for diabetes educators: a companion document to the guidelines for the practice of diabetes education. http://www.diabeteseducator. org/ProfessionalResources/position/competencies.html. Accessed June 26, 2012.
- 90. American Association of Diabetes Educators. A sustainable model of diabetes self-management education/training involves a multi-level team that can include community health workers. http://www.diabeteseducator.org/DiabetesEducation/position/ White\_Papers.html.
- 91. Gillett M, Dallosso HM, Dixon S, et al. Delivering the Diabetes Education and Self Management for Ongoing and Newly Diagnosed (DESMOND) programme for people with newly diagnosed type 2 diabetes: cost effectiveness analysis. *BMJ*. 2010;341:c4093.
- 92. Redman BK. *The Practice of Patient Education*. 10th ed. St. Louis, MO: Mosby; 2007.
- Mulcahy K, Maryniuk M, Peeples M, et al. Diabetes selfmanagement education core outcomes measures. *Diabetes Educ*. 2003;29(5):768-770, 773-784, 787-768.
- Reader D, Splett P, Gunderson EP. Impact of gestational diabetes mellitus nutrition practice guidelines implemented by registered dietitians on pregnancy outcomes. J Am Diet Assoc. 2006;106(9):1426-1433.
- 95. Boucher JL, Evert A, Daly A, et al. American Dietetic Association revised standards of practice and standards of professional performance for registered dietitians (generalist, specialty, and advanced) in diabetes care. J Am Diet Assoc. 2011;111(1): 156-166,e151-e127.
- 96. American Diabetes Association. Standards of medical care in diabetes: 2012. *Diabetes Care*. 2012;35(suppl 1):S11-S63.
- Bantle JP, Wylie-Rosett J, Albright AL, et al. Nutrition recommendations and interventions for diabetes: a position statement of the American Diabetes Association. *Diabetes Care*. 2008;31(suppl 1):S61-S78.
- Wagner EH, Austin BT, Von Korff M. Organizing care for patients with chronic illness. *Milbank Q*. 1996;74(4):511-544.
- 99. Norris SL. Health-related quality of life among adults with diabetes. *Curr Diab Rep.* 2005;5(2):124-130.

- 100. Herman AA. Community health workers and integrated primary health care teams in the 21st century. J Ambul Care Manage. 2011;34(4):354-361.
- 101. Weinger K, Beverly EA, Lee Y, et al. The effect of a structured behavioral intervention on poorly controlled diabetes: a randomized controlled trial. Arch Intern Med. 2011;171(22):1990-1999.
- 102. Norris SL, Zhang X, Avenell A, et al. Long-term effectiveness of lifestyle and behavioral weight loss interventions in adults with type 2 diabetes: a meta-analysis. *Am J Med.* 2004;117(10):762-774.
- 103. Ellis SE, Speroff T, Dittus RS, et al. Diabetes patient education: a meta-analysis and meta-regression. *Patient Educ Couns*. 2004;52(1):97-105.
- 104. Armour TA, Norris SL, Jack L, Jr, et al. The effectiveness of family interventions in people with diabetes mellitus: a systematic review. *Diabet Med.* 2005;22(10):1295-1305.
- 105.Magee M, Bowling A, Copeland J, et al. The ABCs of diabetes: diabetes self-management education program for African Americans affects A1C, lipid-lowering agent prescriptions, and emergency department visits. *Diabetes Educ*. 2011;37(1):95-103.
- 106. Cavanaugh K, Huizinga MM, Wallston KA, et al. Association of numeracy and diabetes control. Ann Intern Med. 2008;148(10):737-746.
- 107. Rothman RL, DeWalt DA, Malone R, et al. Influence of patient literacy on the effectiveness of a primary care-based diabetes disease management program. JAMA. 2004;292(14):1711-1716.
- 108. Schillinger D, Grumbach K, Piette J, et al. Association of health literacy with diabetes outcomes. *JAMA*. 2002;288(4):475-482.
- 109. Rubin RR, Peyrot M, Saudek CD. The effect of a diabetes education program incorporating coping skills, training on emotional well-being, and diabetes self-efficacy. *Diabetes Educ*. 1993;19(3):210-214.
- 110. Trento M, Passera P, Borgo E, et al. A 5-year randomized controlled study of learning, problem solving ability, and quality of life modifications in people with type 2 diabetes managed by group care. *Diabetes Care*. 2004;27(3):670-675.
- 111. Izquierdo RE, Knudson PE, Meyer S, et al. A comparison of diabetes education administered through telemedicine versus in person. *Diabetes Care*. 2003;26(4):1002-1007.
- 112. Garrett N, Hageman CM, Sibley SD, et al. The effectiveness of an interactive small group diabetes intervention in improving knowledge, feeling of control, and behavior. *Health Promot Pract*. 2005;6(3):320-328.
- 113. Piette JD, Glasgow R. Strategies for improving behavioral health outcomes among patients with diabetes: self-management, education. In Gerstein HC, & Haynes RB, eds. *Evidence-Based Diabetes Care*. Ontario, Canada: BC Decker Publishers; 2001:207-251.
- 114.Boren SA. AADE7TM self-care behaviors: systematic reviews. *Diabetes Educ.* 2007;33(6):866, 871.
- 115. American Association of Diabetes Educators. AADE 7 self-care behaviors, American Association of Diabetes Educators position statement. http://www.diabeteseducator.org/DiabetesEducation/ position/position\_statements.html. Accessed June 26, 2012.
- 116. American Association of Diabetes Educators. AADE position statement: individualization of diabetes self-management education. *Diabetes Educ*. 2007;33(1):45-49.
- 117. Gilden JL, Hendryx M, Casia C, et al. The effectiveness of diabetes education programs for older patients and their spouses. J Am Geriatr Soc. 1989;37(11):1023-1030.
- 118.Brown SA. Effects of educational interventions in diabetes care: a meta-analysis of findings. *Nurs Res.* 1988;37(4):223-230.

- 119. Barlow J, Wright C, Sheasby J, et al. Self-management approaches for people with chronic conditions: a review. *Patient Educ Couns*. 2002;48(2):177-187.
- 120. Skinner TC, Cradock S, Arundel F, et al. Four theories and a philosophy: self-management education for individuals newly diagnosed with type 2 diabetes. *Diabetes Spectrum*. 2003;16(2):75-80.
- 121.Brown SA, Hanis CL. Culturally competent diabetes education for Mexican Americans: the Starr County Study. *Diabetes Educ*. 1999;25(2):226-236.
- 122. Sarkisian CA, Brown AF, Norris KC, et al. A systematic review of diabetes self-care interventions for older, African American, or Latino adults. *Diabetes Educ*. 2003;29(3):467-479.
- 123. Chodosh J, Morton SC, Mojica W, et al. Meta-analysis: chronic disease self-management programs for older adults. Ann Intern Med. 2005;143(6):427-438.
- 124. Anderson-Loftin W, Barnett S, Bunn P, et al. Soul food light: culturally competent diabetes education. *Diabetes Educ*. 2005;31(4):555-563.
- 125. Mensing CR, Norris SL. Group education in diabetes: effectiveness and implementation. *Diabetes Spectrum*. 2003;16(2):96-103.
- 126. Brown SA, Blozis SA, Kouzekanani K, et al. Dosage effects of diabetes self-management education for Mexican Americans: the Starr County Border Health Initiative. *Diabetes Care*. 2005;28(3):527-532.
- 127. Hosey GM, Freeman WL, Stracqualursi F, et al. Designing and evaluating diabetes education material for American Indians. *Diabetes Educ.* 1990;16(5):407-414.
- 128. Thomson FJ, Masson EA. Can elderly patients co-operate with routine foot care? *Diabetes Spectrum*. 1995;8:218-219.
- 129. Hawthorne K, Robles Y, Cannings-John R, et al. Culturally appropriate health education for type 2 diabetes in ethnic minority groups: a systematic and narrative review of randomized controlled trials. *Diabet Med.* 2010;27(6):613-623.
- 130. Cavanaugh K, Wallston KA, Gebretsadik T, et al. Addressing literacy and numeracy to improve diabetes care: two randomized controlled trials. *Diabetes Care*. 2009;32(12):2149-2155.
- 131.Doak CC, Doak LG, Root JH. *Teaching Patients with Low Literacy Skills*. Philadelphia, PA: Lippincott; 2008.
- 132. Schillinger D, Piette J, Grumbach K, et al. Closing the loop: physician communication with diabetic patients who have low health literacy. *Arch Intern Med.* 2003;163(1):83-90.
- 133. Channon SJ, Huws-Thomas MV, Rollnick S, et al. A multicenter randomized controlled trial of motivational interviewing in teenagers with diabetes. *Diabetes Care*. 2007;30(6):1390-1395.
- 134. Naik AD, Palmer N, Petersen NJ, et al. Comparative effectiveness of goal setting in diabetes mellitus group clinics: randomized clinical trial. *Arch Intern Med.* 2011;171(5):453-459.
- 135. Glasgow RE, Funnell MM, Bonomi AE, et al. Self-management aspects of the improving chronic illness care breakthrough series: implementation with diabetes and heart failure teams. *Ann Behav Med.* 2002;24(2):80-87.
- 136. Estey AL, Tan MH, Mann K. Follow-up intervention: its effect on compliance behavior to a diabetes regimen. *Diabetes Educ*. 1990;16(4):291-295.
- 137. Beverly EA, Ganda OP, Ritholz MD, et al. Look who's (not) talking: diabetic patients' willingness to discuss self-care with physicians. *Diabetes Care*. 2012;35(7):1466-1472.

- 138. Mulvaney SA, Rothman RL, Wallston KA, et al. An Internetbased program to improve self-management in adolescents with type 1 diabetes. *Diabetes Care*. 2010;33(3):602-604.
- 139.Osborn CY, Mayberry LS, Mulvaney SA, et al. Patient web portals to improve diabetes outcomes: a systematic review. *Curr Diab Rep.* 2010;10(6):422-435.
- 140. Mulvaney SA, Ritterband LM, Bosslet L. Mobile intervention design in diabetes: review and recommendations. *Curr Diab Rep.* 2011;11(6):486-493.
- 141.Polonsky WH, Fisher L, Earles J, et al. Assessing psychosocial distress in diabetes: development of the diabetes distress scale. *Diabetes Care*. 2005;28(3):626-631.
- 142. Davis ED. Role of the diabetes nurse educator in improving patient education. *Diabetes Educ.* 1990;16(1):36-38.
- 143. Glasgow RE, Davis CL, Funnell MM, et al. Implementing practical interventions to support chronic illness self-management. *Jt Comm J Qual Saf.* 2003;29(11):563-574.
- 144. Daly A, Leontos C. Legislation for health care coverage for diabetes self-management training, equipment and supplies: past, present and future. *Diabetes Spectrum*. 1999;12:222-230.
- 145. Grebe SK, Smith RB. Clinical audit and standardised follow up improve quality of documentation in diabetes care. N Z Med J. 1995;108(1006):339-342.
- 146. Aubert RE, Herman WH, Waters J, et al. Nurse case management to improve glycemic control in diabetic patients in a health maintenance organization: a randomized, controlled trial. *Ann Intern Med.* 1998;129(8):605-612.
- 147. Glasgow RE, Peeples M, Skovlund SE. Where is the patient in diabetes performance measures? The case for including patient-centered and self-management measures. *Diabetes Care*. 2008;31(5):1046-1050.
- 148.Beebe CA, Schmitt SS. Engaging patients in education for selfmanagement in an accountable care environment. *Clin Diabetes*. 2011;29(3):123-126.
- 149. American Association of Diabetes Educators. Standards for outcomes measurement of diabetes self-management education. http://www.diabeteseducator.org/ProfessionalResources/position/ position\_statements.html. Accessed June 26, 2012.
- 150. American Association of Diabetes Educators. Standards for outcomes measurement of diabetes self-management education, technical review. http://www.diabeteseducator.org/ ProfessionalResources/position/position\_statements.html. Accessed June 26, 2012.
- 151. Anderson D, Christison-Lagay J. Diabetes self-management in a community health center: improving health behaviors and clinical outcomes for undeserved patients. *Clin Diabetes*. 2008;26(1):22-27.
- 152. Duncan I, Ahmed T, Li QE, et al. Assessing the value of the diabetes educator. *Diabetes Educ.* 2011;37(5):638-657.
- 153.Institute for Healthcare Improvement. Science of improvement: how to improve. http://www.ihi.org/knowledge/Pages/ HowtoImprove/ScienceofImprovementHowtoImprove.aspx. Accessed June 25, 2012.
- 154. Joint Commission on Accreditation of Healthcare Organizations. Joint Commission Resources: Cost-Effective Performance Improvement in Ambulatory Care. Oakbrook Terrace, IL: Joint Commission on Accreditation of Healthcare Organizations; 2003.

For reprints and permission queries, please visit SAGE's Web site at http://www.sagepub.com/journalsPermissions.nav.

Pages 630-634 was intentionally removed from this PDF. In the print version, these pages were advertisements.